

- 52.** Agostoni C, Axelsson I, Goulet O, Koletzko B, Michaelsen KF, Puntis JW, et al. ESPGHAN Committee on Nutrition. Prebiotic oligosaccharides in dietary products for infants: a commentary by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr.* 2004;39:465-73.
- 53.** Brunser O, Gotteland M, Cruchet S, Figueroa G, Garrido D, Steenhout P. Effect of milk formula with prebiotics on the intestinal microbiota of infants after an antibiotic treatment. *Pediatr Res.* 2006;59:451-6.
- 54.** Moro G, Arslanoglu S, Stahl B, Jelinek J, Wahn U, Boehm G. A mixture of prebiotic oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child.* 2006;91:814-9.
- 55.** Buzzese E, Volpicelli M, Salvini F, Bisceglia M, Lionetti P, Conquetti M, et al. Effect of early administration of GOS/FOS on the prevention of intestinal and extra-intestinal infections in healthy infants. *J Pediatr Gastroenterol Nutr.* 2006;42:e95.
- 56.** Mihatsch WA, Hoegel J, Pohlandt F. Prebiotic oligosaccharides reduce stool viscosity and accelerate gastrointestinal transport in preterm infants. *Acta Paediatr.* 2006;95:843-8.
- 57.** Eiwegger T, Stahl B, Schmitt J, Boehm G, Gerstmayr M, Pichler J, et al. Human milk-derived oligosaccharides stimulate cytokine production of cord blood T-cells in vitro. *Pediatr Res.* 2004;56:536-40.
- 58.** Schelzle H, Wirth S, Skopnik H, Radke M, Knol J, Böckler HM, et al. Randomized double-blind study of the nutritional efficacy and bifidogenicity of a new infant formula containing partially hydrolyzed protein, a high β-palmitic acid level, and nondigestible oligosaccharides. *J Pediatr Gastroenterol Nutr.* 2003;36:343-51.
- 59.** Savino F, Cresi F, Maccario S, Cavallo F, Dalmasso P, Fanaro S, et al. Minor feeding problems during the first months of life: effect of a partially hydrolyzed milk formula containing fructo- and galacto-oligosaccharides. *Acta Paediatr.* 2003;441:S86-90.
- 60.** Scholtens PA, Alles MS, Bindels JG, Van der Linde EG, Tolboom JJ, Knol J. Bifidogenic effects of solid weaning foods with added prebiotic oligosaccharides: a randomized controlled clinical trial. *J Pediatr Gastroenterol Nutr.* 2006;42:553-9.
- 61.** Moore N, Chao C, Yang LP, Oliva-Hemker M, Saavedra JM. Effects of fructo-oligosaccharide-supplemented infant cereal: a double-blind, randomized trial. *Br J Nutr.* 2003;90:581-7.
- 62.** Veereman-Wauters G. Application of prebiotics in infants foods. *Br J Nutr.* 2005;93 Suppl 1:S57-60.
- 63.** Duggan C, Penny M, Hibberd P, Gil A, Huapaya A, Cooper A, et al. Oligofructose-supplemented infant cereal. 2. Randomized, blinded, community-based trials in Peruvian infants. *Am J Clin Nutr.* 2003;77:937-42.
- 64.** Hockstra JH, Szajewska H, Zikri MA, Micetic-Turk D, Weizman Z, Papadopoulou A, et al. Oral rehydration solution containing a mixture of non-digestible carbohydrates in the treatment of acute diarrhea: a multicenter randomized placebo controlled study on behalf of the ESPGHAN Working Group on Intestinal Infections. *J Pediatr Gastroenterol Nutr.* 2004;39:239-45.

Fe de errores

En el artículo “Consenso sobre tratamiento del asma en pediatría”, del Grupo de trabajo sobre Tratamiento del Asma Infantil, publicado en An Pediatr (Barc). 2007;67(3):253-73, se han detectado los siguientes errores:

- Tabla 10 (p. 262): donde dice “GCI dosis medias + ARLT” debe decir “GCI dosis bajas + ARLT”.
- Figura 3 (p. 262): en el paso 5, donde dice “GCI dosis medias + ARLT” debe decir “GCI dosis altas + ARLT”.
- Figura 4 (p. 264): en el paso 4, donde dice “GCI dosis baja + ARLT” debe decir “GCI dosis media + ARLT”; en el paso 5 donde dice “GCI dosis medias” debe decir “GCI dosis altas”; en el paso 6, donde dice “Omalizumab” debe decir “Anticuerpos anti-IgE”.