

## IMAGES IN PAEDIATRICS

**Lice: Be careful, not just on the head!****Pediculosis: ¡atención, no sólo en la cabeza!**


Aniza Giacaman\*, Francisca Mestre-Bauzá

Departamento de Dermatología, Hospital Universitario Son Espases. Palma de Mallorca, Spain

Available online 1 July 2025

A boy aged 4 years presented with bluish-brown spots (maculae ceruleae) on the trunk (Fig. 1A). Lice and nits could be seen on the eye lashes (Fig. 1B). Dermoscopy revealed the presence of crab lice on the scalp, trunk (Fig. 1C) and eye lashes (Fig. 1D). It also allowed visualization of nits, both viable (Fig. 1E) and hatched (Fig. 1F). Sexual abuse was ruled out through a thorough history-taking, with questioning of both child and mother. The physical examination was otherwise normal. We prescribed oral ivermectin to the entire household. We recommended repetition of treatment one week after as well as hygiene measures in the house.

*Pediculus humanus capitis*, *P. humanus corporis* and *Phthirus pubis* are distributed worldwide. The body of *P. pubis* is smaller ( $1.2 \times 0.8$  mm) compared to *P. capitis*, and is brown in color, with an appearance similar to that of a crab, has six legs and clings to hair shafts with the claws of its front legs. Viable nits are oval structures, brown in color, with convex ends. Hatched nits appear see-through and are missing one of the ends. In children, the presence of *P. pubis* should always trigger suspicion of sexual abuse, although the parasite can also spread in crowded conditions through non-genital body contact.<sup>1-3</sup> Maculae ceruleae are associated with pediculosis, do not require treatment and resolve spontaneously when the infestation is treated.<sup>3</sup>



**Figure 1** (a) Maculae ceruleae on the trunk of a boy with a body louse infestation. (b) Lice and nits on the eye lashes. (c) *Phthirus pubis* clinging to hair in the trunk. (d) the same parasite clinging to the patient's eye lashes. (e) Viable nits (black square box). (f) Nonviable or hatched nits.

DOI of original article:

<https://doi.org/10.1016/j.anpedi.2025.503810>

\* Corresponding author.

E-mail address: [anizagiacaman@gmail.com](mailto:anizagiacaman@gmail.com) (A. Giacaman).

## References

1. Salavastru CM, Chosidow O, Janier M, Tiplica GS. European guideline for the management of pediculosis pubis. J Eur Acad Dermatol Venereol. 2017;31:1425–8.
2. Yao B, Hu C, Yue X, Liu G, Wang B. Clinical characteristics and prognosis of pediatric *Phthirus pubis* coinfection of the eyelashes and scalp hairs. Am J Trop Med Hyg. 2023;109:686–9.
3. Haitz K, Walls AC, Huang JT. A 15-month-old boy with blue-gray macules. JAMA Pediatr. 2017;171:191–2. <http://dx.doi.org/10.1001/jamapediatrics.2016.2832>.