



SPANISH ASSOCIATION OF PAEDIATRICS

Use of sugars and sweeteners in children's diets. Recommendations of the Nutrition Committee of the Spanish Association of Paediatrics[☆]



M. Gil-Campos^{a,*}, M.A. San José González^b, J.J. Díaz Martín^c, Comité de Nutrición de la Asociación Española de Pediatría[◇]

^a *Unidad de Metabolismo e Investigación Pediátrica, Hospital Universitario Reina Sofía, Universidad de Córdoba, IMIBIC, Córdoba, Spain*

^b *Atención Primaria, Centro de Salud de Sarria, Lugo, Spain*

^c *Sección de Gastroenterología y Nutrición Pediátrica, Hospital Universitario Central de Asturias, Oviedo, Asturias, Spain*

Received 29 December 2014; accepted 18 February 2015
Available online 11 November 2015

KEYWORDS

Sweeteners;
Children;
Sugars;
Obesity;
Tooth decay

Abstract The term "sweetener" refers to a food additive that imparts a sweet flavour and usually provides no or very low energy. It is used to sweeten foods, medicines and food supplements with no nutritional purposes. For years, no-calorie sweeteners have been used as substitutes for all or part of the sugar content in foods and beverages. In recent decades its consumption has risen to prevent tooth decay, or as an aid in weight control, obesity and diabetes and, in general, to achieve an optimal energy balance. However, consumption of sugary or sweetened food and soft drinks is high, making this situation of special interest in calorie intake and in the poor behavioural pattern of eating habits in children. In addition, questions remain among consumers about the risks to health associated with their use, whether they are artificial or natural. The "artificial sweeteners" are the group of greatest interest in research in order to demonstrate their safety and to provide firm data on their possible therapeutic effects. The aim of the present document is to increase information for paediatricians on the characteristics of different sweeteners, and to advise on the choice of sweeteners, based on their properties.

© 2014 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. All rights reserved.

[☆] Please cite this article as: Gil-Campos M, San José González MA, Díaz Martín JJ. Uso de azúcares y edulcorantes en la alimentación del niño. Recomendaciones del Comité de Nutrición de la Asociación Española de Pediatría. An Pediatr (Barc). 2015;83:353.e1–353.e7.

* Corresponding author.

E-mail address: mercedes_gil_campos@yahoo.es (M. Gil-Campos).

◇ The members of the Comité de Nutrición de la AEP are presented in [Appendix A](#).

PALABRAS CLAVE

Edulcorantes;
 Infancia;
 Azúcares;
 Obesidad;
 Caries

Uso de azúcares y edulcorantes en la alimentación del niño. Recomendaciones del Comité de Nutrición de la Asociación Española de Pediatría

Resumen El término edulcorante hace referencia a aquel aditivo alimentario que confiere un sabor dulce y que, habitualmente, no aporta o proporciona muy poca energía. Se utiliza para endulzar alimentos, medicamentos y complementos alimenticios cuando se persiguen fines no nutritivos. Desde hace años, se han empleado edulcorantes acalóricos como sustitutos de todo o parte del contenido en azúcares en comidas y bebidas. En las últimas décadas, se ha incrementado su consumo para prevenir la caries y para el correcto cumplimiento de la dieta en casos de control del peso corporal, obesidad y diabetes y, en general, como coadyuvantes para conseguir un balance energético adecuado. No obstante, el consumo de alimentos y de bebidas azucaradas y/o edulcoradas es elevado, reflejando o un aporte calórico importante, o un patrón de hábitos alimentarios inadecuados en los niños. Por otro lado, sigue habiendo dudas entre los consumidores sobre los riesgos para la salud asociados al uso de edulcorantes, ya sean artificiales o naturales. El principal interés en investigación sobre seguridad y los posibles usos terapéuticos se centra en los «edulcorantes artificiales». El objetivo de este documento es proporcionar información a los pediatras sobre las características de los distintos edulcorantes para aconsejar en la elección de un determinado edulcorante sobre la base de sus propiedades. © 2014 Asociación Española de Pediatría. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Sugars and sweeteners: concepts

Carbohydrates (CHs) are the nutrients that constitute the main source of food energy. They are notable for their structure and pleasant taste, which in some cases, such as sugars, is sweet, making other foods more palatable, for their ability to satiate the appetite and, in some, for their high fibre content.¹ CHs should provide between 45% and 60% of total dietary energy intake in children older than one year of age.² CHs from food are presented in the form of complex molecules (polymers or polysaccharides), especially starches, dextrans and fibre, or simpler ones, commonly called sugars. The main dietary sources of sugars are fruit and fruit juices, some vegetables, milk, and processed foods with added sugars, especially sucrose or hydrolysed starch (glucose or fructose syrups), such as soft drinks, pastries, sweets and confectionery.^{1,2} Sugars are used to sweeten or enhance the flavour of many of them, to modify the freezing and melting point or to colour foods naturally, and to preserve them. Balanced intake of sugars in the daily diet has important properties, as it facilitates the rapid supply of glucose, an indispensable carbohydrate for the development of cognitive functions and physical activity, to the brain and muscles. Sugar should be consumed in a natural form with the foods that contain it, since this also provides other micronutrients. In the twentieth century, however, questions began to be raised as to whether excessive consumption of sugars, particularly associated with processed foods, might be related to diabetes or obesity, and research has continued up to the present.³⁻⁵

Additives are substances deliberately added to foods to perform certain technological functions and the result is that both the additive itself and its byproducts become components of those foods. Additives are not consumed as foods or used as typical ingredients in the diet, regardless of whether or not they have nutritional value.

Monosaccharides, disaccharides or oligosaccharides, or foods containing them, are not regarded as food additives. The term *sweetener* refers to a substance used to impart a sweet taste to foods (Regulation (EC) No 1333/2008). Thus foods such as honey or ordinary sugar, fructose or glucose are not considered sweeteners, since they have other functions apart from sweetening.⁶

Low- or no-calorie sweeteners (LNCSs) have been used for years to replace all or part of the sugar content in foods and drinks, but in the last few decades their consumption has increased both in adults and in children. Their use is linked to dietary alternatives for weight control or diabetes, but also to preventing tooth decay. Although some studies question these possible benefits, both in adults and in children,^{7,8} systematic reviews and meta-analyses on this subject conclude that the use of sweeteners is beneficial in weight-control and diabetes programmes associated with a healthy lifestyle.⁹

The purpose of this document is to provide paediatricians with information on sweeteners and health-related issues, in order to give appropriate advice to patients and their families.

Consumption of sugars and sweeteners and its relationship with health

To assess this consumption we have to take into account not only added sugars and sweeteners, but also sugar incorporated as an ingredient in precooked/processed foods. European adolescents consume some 384 kcal per day from drinks, of which 30.4%, 20.7% and 18.1% comes from sugar-sweetened beverages, sweetened milk and fruit juice, respectively.¹⁰ Various cross-sectional studies have concluded that there is no association, or even that there is a negative association, between consumption of sugars and weight gain.^{11,12} However, there is a widespread debate on

whether greater intake of sugars through sweetened beverages could have a significant effect on increase in body mass index (BMI) or about whether unsweetened diets may have an influence on control of obesity.¹³

Some observational studies have also related increase in BMI to consumption of non-caloric sweeteners, although these data should not be interpreted as proof of a causal relationship, but rather as a sign that the probability of consumption is higher in the obese and sedentary population (reverse causality).⁸ Replacing sugar with sweeteners is not necessarily associated with a lower overall calorie intake, and could encourage unbalanced dietary habits, involving high consumption of low-calorie products with sweeteners and of others with excessive calories.^{6,14} Some authors postulate that early consumption of sugar-sweetened products in infants and young children could affect self-regulation of eating and preference for sweet flavours,^{7,15} which may be maintained during childhood and adolescence. It has even been suggested that consuming sweeteners and sugars together in the diet could give rise to a neuronal response that leads to more rapid absorption of sugars, also increasing secretion of peptides related to glucagon, or insulin.¹⁶ On this subject a highly controversial hypothesis has recently been put forward on the development of glucose intolerance through alterations to the intestinal microbiota in mice associated with consuming saccharin at maximum doses.¹⁷ However, one must be very cautious in extrapolating conclusions from animal studies to humans.¹⁸ Nowadays diabetic patients can use non-caloric sweeteners as part of a balanced, controlled diet, and studies in adults indicate that they do not affect plasma glucose or lipid levels, although this has not been sufficiently investigated in children.^{19,20}

Fermentable CHs in the diet produce acidic materials which destroy hard tissues of the tooth, giving rise to tooth decay. In the child population dental caries is still common, because of a general lack of adequate oral hygiene after consuming sugary foods, as well as fluoride deficiency.²¹ In infants, moreover, prolonged use of fruit juices or other fructose- or sucrose-rich sweetened beverages in dummies increases the risk of damage to teeth. Among sweeteners, polyols (sugar alcohols) and sucralose, among others, have been declared to have healthy properties for avoiding this condition.²²

Recommendations for consumption

All types of CHs must be represented in a healthy diet. The FAO and WHO recommend an intake of sugars (simple carbohydrates) of less than 10% of the total caloric value of the diet, trying to make them part of a healthy diet with limited consumption of sugar-sweetened drinks.²³ In 2005 it was established that the recommended daily intake in adults and children older than one year of age is 100 g/day of CHs as an estimated mean requirement.²⁴ In Spain, adhering to a Mediterranean or similar diet would make it possible to maintain an appropriate energy intake without sacrificing its sweetening and pleasurable function. In addition, as a practical measure to achieve greater benefits, it has been recommended that consumption of sugars or sugar-sweetened foods should be limited to less than 3 times per

day, not exceeding 6% of total energy intake,²⁵ and that of sugar-sweetened drinks to occasional use. It is also recommended that excessive consumption of high-fructose corn syrup (55% fructose) should be reduced, as it represents a health hazard, especially for children.²⁶ Indeed, these syrups are not used in Spain, where the maximum fructose content of manufactured products is approximately 50%.

The nutrition statements prescribed in Regulation (EC) No 1924/2006 make it easy to ascertain the composition of foods in terms of sugars. Thus according to the labelling we can find foods that are "low in sugar" (no more than 5 g of sugars per 100 g for solids or 2.5 g of sugars per 100 ml for liquids), "sugar-free" (no more than 0.25 g per 100 g or 100 ml), or "with no added sugar" (a product with no added mono- or disaccharides nor any other food used for its sweetening properties). In the last of these cases, the food may "contain naturally occurring sugars" or "sugar(s) and sweetener(s)", and this information must appear on the nutrition information label.²⁷ On a practical level, the American Heart Association has suggested simple advice, such as not having more than 6 teaspoonfuls of sugar per day for women (25 g) (e.g., present in 250 ml of a sugar-sweetened beverage) and 9 for men (37 g).²⁸

During pregnancy CH intake should not be restricted, but intake of refined sugar or foods with added sugar should be controlled. There is an increase in availability of glucose, which is an essential substrate for the foetus. In newborns, because of pancreatic amylase deficiency, it is advisable for infant milk formulae not to contain starch, with lactose as the main CH, as well as glucose polymers, which can also be added, since they are directly absorbable by the enterocyte. On the other hand, the infant's predisposition towards sweet taste makes it possible to introduce new foods with a relatively high starch content.²⁹

As for sweeteners, the European Food Safety Authority (EFSA) proposes acceptable daily intakes to ensure appropriate use without possible adverse effects.³⁰ In the case of infants it is difficult to find sweeteners in the products they consume, as they are prohibited in milk formulae, cereals and baby foods. At later ages, foods such as soft drinks, fruit juices or dairy products sweetened with sugar or sweeteners should be controlled. Reducing the energy content in some foods through sweeteners does actually seem to have modest benefits in children.¹⁶ On the other hand, it may lead to greater consumption of other more caloric foods, and there is also the possibility that early exposure to sweeteners, as mentioned previously, could affect dietary habits later.³¹ In pre-school and school-aged children the recommendations already described should be maintained, taking some additional factors into account, such as physical activity, which may mean that sugar intake has to be regulated with appropriate beverages rather than just drinking water.³² Indeed, anaerobic or prolonged exercise causes a depletion of liver and muscle glycogen reserves, leading to fatigue and cognitive abnormalities,³³ which sometimes have to be mitigated with sugars.

Types of sweeteners

There are various ways of classifying sweeteners (Tables 1 and 2). Using the glycaemic index (GI) we

Table 1 Classification of sugars and sweeteners.

Natural	Caloric	Sugars	Sucrose, glucose, galactose, fructose, lactose, maltose, trehalose
		Others	Honey, maple syrup, palm sugar, coconut sugar
	Non-caloric	Luo han guo, stevia, thaumatin, pentadin, monellin, brazzein	
Artificial	Caloric	Modified sugars	(High fructose) corn syrup, inverted sugar
		Sugar alcohols	Sorbitol, xylitol, mannitol, lactitol
	Non-caloric	Artificial sweeteners	Aspartame, sucralose, saccharin, neotame, acesulfame K, cyclamic acid, alitame, advantame

Source: Modified from García-Almeida et al.³⁷

can also make a classification of foods based on postprandial blood sugar response, comparing them with a reference food (GI = 100). Sucrose is medium GI (≈ 65).³⁴ Depending on their energy contribution they are classified as “caloric” or “low calorie/non-caloric” (Table 1) or as “nutritive” or “non-nutritive”, as established by the US Food and Drug Administration (FDA) (Table 2).³⁵ In some cases the term “natural origin” is used in the consumer information, as with steviol glycoside, or “artificial” origin when they are synthetic (Table 1).

Natural caloric sweeteners include sucrose, fructose, glucose and maltose. Fructose has classically been used as a substitute for sucrose in diabetic patients. More recently, however, it has been found that high-fructose diets, especially if the fructose is added to processed foods, could induce hyperinsulinaemia, hypertriglyceridaemia and insulin resistance, and this has led to a recommendation to limit their use in diabetics. In addition, fructose dependence has been described, leading to high consumption.³⁶

Fructooligosaccharides have a sweetening power relative to sucrose of 0.3–0.6. Inulin is a fructan with a well-known prebiotic effect found naturally in the yacon or Peruvian ground apple, a tuber native to the Andes, considered a

Table 2 Classification of sweeteners according to the US Food and Drug Administration (FDA) and the European Union, and the sweetening power (SP) of each substance relative to sucrose (table sugar): degree of sweetness comparable with sugar (bulk sweetener) or much higher (intense sweetener).

Sweeteners	Classification by caloric contribution ^a	Classification by sweetening power ^b
<i>Polyols</i>		Bulk
Sorbitol E-420	Nutritive	0.5–1
Mannitol E-421	Nutritive	0.7
Isomalt E-953	Nutritive	0.45–0.65
Maltitol E-965	Nutritive	1
Lactitol E-966	Nutritive	0.5
Xylitol E-967	Nutritive	1
Erythritol E-968	Non-nutritive	0.7
<i>Acesulfame K E-950</i>	Non-nutritive	Intense (200)
<i>Aspartame E-951</i>	Nutritive	Intense (150–200)
<i>Cyclamic acid E-952</i>	Non-nutritive	Intense (50–100)
<i>Saccharin E-954</i>	Non-nutritive	Intense (300–400)
<i>Thaumatin E-957</i>	Nutritive	Intense (2500)
<i>Neohesperidin DC E-959</i>	Nutritive	Intense (250–2000)
<i>Neotame E-961</i>	Non-nutritive	Intense (7000–13,000)
<i>Sucralose E-955</i>	Non-nutritive	Intense (600)
<i>Steviol glycoside E-960</i>	Non-nutritive	Intense (350–400)

^a In intense sweeteners the energy contribution to the diet is insignificant, because the amount of product consumed is very small.

^b In parentheses, sweetening power value relative to sucrose (value 1).

health food on account of being rich in various minerals, vitamin C and B-group vitamins. Coconut sugar is another traditional product with a low GI because of its sucrose content. Polyols derived from sugar and also regarded as CHs are produced, albeit in small quantities, in plants and cereals. They generally contain fewer calories than sugar, with a very low GI, and have not been associated with the development of tooth decay, since they are non-fermentable; they are used as sweeteners. Their intestinal absorption is generally low, and in moderate quantities (more than 10 g of sorbitol) they can produce flatulence, colic or diarrhoea, as they are fermented by the colonic microbiota.

Industrial processed foods use sugars converted from starch, with a high GI and high calorie content. Corn syrup is the prime example.

There are also natural sweeteners (stevia, luohanguo, thaumatin and brazzein) with no significant calorie content in the quantities normally consumed for sweetening purposes. These are not CHs, and therefore have no GI. They are regarded as high-intense sweeteners.

‘Artificial sweeteners’ are characterised by being non-caloric, with no glycaemic effect and high sweetening power (Table 2). Prominent among them is saccharin, which has great sweetening capacity. Sucralose and aspartame are also notable for their extensive worldwide use, especially in beverages.³⁷ Aspartame is made from a methyl ester of phenylalanine and aspartic acid. Its use is approved by the FDA and the EFSA.³⁸

Safe use of sweeteners

Society demands a market supply of high-quality food substances that are suitable for consumption by people with specific needs, such as diabetics, or that respond to the current demand for low-calorie products. Although the object of using LNCSs is to reduce calorie intake in the diet and the presence of fermentable sugars in the mouth, seeking beneficial effects, such as promoting a decrease in body weight, and preventing diabetes or tooth decay, consumers have doubts about the health risks associated with their use, depending on whether they are artificial or natural. The fact that a sweetener is of natural origin does not imply greater safety or effectiveness. Since a wide range of molecules is involved, there are many potential sources of risk: interference in absorption, metabolism or excretion of nutrients or intermediate metabolites, as well as allergic reactions, accumulation in tissues, effects on normal intestinal flora, disruption of blood glucose regulation or interaction with other drugs or medications.

In research on safety and possible therapeutic uses in patients with diabetes or other specific health problems the focus of interest is on ‘artificial sweeteners’. For this purpose the legal aspects of safety and efficacy in the use of sweeteners need to be constantly reviewed by the EFSA.

Regulation (EC) No 1333/2008 lays down rules on the food additives used in foods to ensure the protection of consumers' health while upholding trade practices. Annex II, updated in 2013, indicates that the following LNCSs are authorised: acesulfame-K (E-950), aspartame (E-951), salt of aspartame-acesulfame (E-962), cyclamic acid and its Na and Ca salts (E-952), neohesperidine DC (E-959), saccharin and its Na, K and Ca salts (E-954), sucralose (E-955), thaumatin (E-957) and neotame (E-961). Following the favourable verdict from the EFSA in 2010, the use of stevia derivatives, steviol glycosides (E-960), as natural non-caloric sweeteners was definitively approved throughout the European market, and also sorbitol and xylitol. In 2014 the use of advantame (E-969) as an intense sweetener was authorised.

With regard to aspartame, it does not represent a risk of toxicity for consumers at current exposure levels. Because it is a source of phenylalanine it is not recommended in individuals with hyperphenylalaninaemia or phenylketonuria, although in studies in patients with consumption at normal doses and long-term studies, even in pregnant women, no adverse effects have been found.³⁷

Information for correct use of these substances is based on knowing the differences between the information labels of commonly consumed products containing sweeteners.

The labelling of food additives must comply with the general conditions laid down in Directive 2000/13/EC. Specifically, if the product contains polyols, the label must state: ‘excessive consumption may produce laxative effects’, or if it contains aspartame or salt of aspartame-acesulfame: ‘contains a source of phenylalanine’. In this regard, however, there is substantial room for improvement to ensure that this compulsory information appears on all products containing sweeteners.

Under European regulations (Council Directive 89/398/EEC), artificial sweeteners must not be used in infant formulae, follow-on formulae, cereals, baby foods or dietary foods for very young children for special medical purposes, unless expressly indicated.³⁸ Sugars such as sucrose or fructose may be added in limited quantities.

Scientific research, though still scarce in humans, shows that sweeteners are safe in the general population, including pregnant women³⁹ and children, although in these populations they must be used in moderation.^{5,16,22}

All the food additives approved in the European Union are considered safe in the specified doses and conditions of use. In children, however, additives of this kind should only be used as an alternative when other forms of obesity prevention are not sufficient, except for the use of sugar-free chewing gum to prevent tooth decay, or use in pharmaceutical products.⁵

Final comments

1. In general, an intake of simple CHs of less than 10% of the caloric value of the diet is appropriate in the context of a healthy lifestyle.
2. Sweeteners, especially ‘non-caloric’ sweeteners, can help to limit the intake of refined sugars in the diet and are useful for preventing diseases such as obesity and diabetes, in association with a moderate, balanced diet.
3. In view of the recommendation not to add sweeteners to foods intended for infants and very young children, their use is not advisable in children between the ages of 1 and 3 years.
4. It is recommended that health care professionals acquire the appropriate knowledge to advise on and/or choose a particular sweetener on the basis of its properties.
5. There is a continuing need for specific research in order to make appropriate use of sweeteners in children.

Conflicts of interest

The authors have no conflicts of interest to declare.

Appendix A. The members of the Nutrition Committee of the Spanish Association of Paediatrics

José Manuel Moreno Villares (coordinator), Juan José Díaz Martín, Mercedes Gil Campos, Ana Moráis López, Víctor Manuel Navas López, Susana Redecillas Ferreiro, Miguel

Sáenz de Pipaón, Miguel Ángel San José González, Félix Sánchez-Valverde Visus.

References

1. FAO/WHO (Food and Agriculture Organization/World Health Organization). Carbohydrates in human nutrition. (FAO Food and Nutrition Paper-66). Rome: FAO; 1998.
2. EFSA Panel on Dietetic Products, Nutrition, and Allergies (NDA). Scientific opinion on dietary reference values for carbohydrates and dietary fibre. *EFSA J.* 2010;8:1462–77.
3. Food National Board. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (2002/2005). A report of the Panel on Macronutrients, Subcommittees on Upper Reference Levels of Nutrients and Interpretation and Uses of Dietary Reference Intakes, and the Standing Committee on the Scientific Evaluation of Dietary Reference Intakes; 2005. Available from: <http://www.nap.edu/openbook> [accessed 11.12.14].
4. Welsh JA, Cunningham SA. The role of added sugars in pediatric obesity. *Pediatr Clin N Am.* 2011;58:1455–66.
5. Serra-Majem L, Riobó P, Belmonte S, Anadón A, Aranceta J, Franco E, et al. Chinchón declaration; decalogue on low- and no-calorie sweeteners (LNCS). *Nutr Hosp.* 2014;29:719–34.
6. Gómez Candela C, Palma Milla S. Una visión global, actualizada y crítica del papel del azúcar en nuestra alimentación. *Nutr Hosp.* 2013;28 Suppl. 4:1–4.
7. Fernstrom JD, Navia JL. Supplement: low-calorie sweeteners and weight control – what the science tells us. *J Nutr.* 2012;142:1170S–2S.
8. Foreyt J, Kleinman R, Brown RJ, Lindstrom R. The use of low-calorie sweeteners by children: implications for weight management. *J Nutr.* 2012;142:1156S–62S.
9. Miller PE, Perez V. Low-calorie sweeteners and body weight and composition: a meta-analysis of randomized controlled trials and prospective cohort studies. *Am J Clin Nutr.* 2014;100:765–77.
10. Duffey KJ, Huybrecht I, Mouratidou T, Libuda L, Kersting M, DeVriendt T, et al. Beverage consumption among European adolescents in the HELENA Study. *Eur J Clin Nutr.* 2012;66:244–52.
11. Saris WH, Astrup A, Prentice AM, Zunft HJ, Formiguera X, Verboeket-van de Venne WP, et al. Randomized controlled trial of changes in dietary carbohydrate/fat ratio and simple vs. complex carbohydrates on body weight and blood lipids: the CARMEN study. The carbohydrate ratio management in European national diets. *J Obes Relat Metab Disord.* 2000;24:1308–10.
12. Barclay AW, Brand-Miller J. The Australian paradox: a substantial decline in sugars intake over the same timeframe that overweight and obesity have increased. *Nutrients.* 2011;3:491–504.
13. Yang Q. Gain weight by going diet? Artificial sweeteners and the neurobiology of sugar cravings: Neuroscience 2010. *Yale J Biol Med.* 2010;83:101–8.
14. Wiebe N, Padwal R, Field C, Marks S, Jacobs R, Tonelli M. A systematic review on the effect of sweeteners on glycemic response and clinically relevant outcomes. *BMC Med.* 2011;9:123.
15. Drewnowski A1, Mennella JA, Johnson SL, Bellisle F. Sweetness and food preference. *J Nutr.* 2012;142:1142S–8S.
16. Brown RJ, de Banate MA, Rother KI. Artificial sweeteners: a systematic review of metabolic effects in youth. *Int J Pediatr Obes.* 2010;5:305–12.
17. Suez J, Korem T, Zeevi D, Zilberman-Schapira G, Thaiss CA, Maza O, et al. Artificial sweeteners induce glucose intolerance by altering the gut microbiota. *Nature.* 2014;514:181–6.
18. Calorie Control Council: Comments on artificial sweeteners induce glucose intolerance by altering the gut microbiota. *Nature.* 2014;514:181–6.
19. Scientific Opinion on the substantiation of health claims related to intense sweeteners and contribution to the maintenance or achievement of a normal body weight (ID 1136, 1444, 4299), reduction of post-prandial glycaemic responses (ID 4298), maintenance of normal blood glucose concentrations (ID 1221, 4298), and maintenance of tooth mineralisation by decreasing tooth demineralisation (ID 1134, 1167, 1283) pursuant to Article 3(1) of Regulation (EC) No 1924/2006. *EFSA J.* 2011;9:2226.
20. Raben A, Richelsen B. Artificial sweeteners: a place in the field of functional foods? Focus on obesity and related metabolic disorders. *Curr Opin Clin Nutr Metab Care.* 2012;15:597–604.
21. Anderson CA, Curzon ME, van Loveren C, Tatsi C, Duggal MS. Sucrose and dental caries: a review of the evidence. *Obes Rev.* 2009;10 Suppl. 1:41–54.
22. Fitch C, Keim KS, Academy of Nutrition and Dietetics. Position of the Academy of Nutrition and Dietetics: use of nutritive and nonnutritive sweeteners. *J Acad Nutr Diet.* 2012;112:739–58.
23. OMS/FAO. Dieta, nutrición y prevención de enfermedades crónicas de 2003. Ginebra: Informe Técnico. OMS; 2003. Available from: http://whqlibdoc.who.int/trs/WHO_TRS_916_spa.pdf [accessed 05.12.14].
24. Organización de las Naciones Unidas para la Alimentación y la Agricultura (FAO). Hojas de balance alimentario; 2009. Available from: <http://faostat.fao.org/site/368/default.aspx#ancor> [accessed 04.11.14].
25. Objetivos poblacionales para la población española. Consenso de la Sociedad Española de Nutrición Comunitaria, 2011. *Rev Esp Nutr Comun.* 2011;17:178–99.
26. Simopoulos AP, Bourne PG, Faergeman O. Bellagio report on healthy agriculture, healthy nutrition, healthy people. *Nutrients.* 2013;5:411–23.
27. Reglamento (CE) 1924/2006, de 20 de diciembre, relativo a las declaraciones nutricionales y de propiedades saludables en los alimentos. DOCE de 30/12/2006.
28. Johnson RK, Appel LJ, Brands M, Howard BV, Lefevre M, Lustig RH, et al. Dietary sugars intake and cardiovascular health: a scientific statement from the American Heart Association. *Circulation.* 2009;120:1011–20.
29. Partearroyo T, Sánchez Campayo E, Varela Moreiras G. El azúcar en los distintos ciclos de la vida: desde la infancia hasta la vejez. *Nutr Hosp.* 2013;28 Suppl. 4:40–7.
30. Guidance for submission for food additive evaluations. *EFSA J.* 2012;10:2760.
31. Sylvestry A, Rother KI, Brown R. Artificial sweetener use among children: epidemiology, recommendations, metabolic outcomes, and future directions. *Pediatr Clin N Am.* 2011;58:1467–80.
32. Sünram-Lea SI, Foster JK, Durlach P, Pérez C. Glucose facilitation of cognitive performance in healthy young adults: examination of the influence of fast-duration, time of day and preconsumption plasma glucose levels. *Psychopharmacology (Berl).* 2001;157:46–54.
33. Rennie KL, Livingstone MB. Associations between dietary added sugar intake and micronutrient intake: a systematic review. *Br J Nutr.* 2007;97:832–41.
34. Brownell K, Gold M, editors. Handbook of food and addiction. New York: Oxford University Press; 2012.
35. Position of American Dietetic Association: use of nutritive and nonnutritive sweeteners. *J Am Diet Assoc.* 2004;104:255–75.
36. Atkinson FS, Foster-Powell K, Brand-Miller JC. International tables of glycemic index and glycemic load values: 2008. *Diabetes Care.* 2008;31:2281–3.

37. García-Almeida JM, Casado Fdez GM, García Alemán J. Una visión global y actual de los edulcorantes. Aspectos de regulación. *Nutr Hosp.* 2013;28 Suppl. 4:17–31.
38. Directiva 94/35/CE del Parlamento Europeo y del Consejo, de 30 de junio de 1994. Diario Oficial de las Comunidades Europeas. Available from: http://europa.eu/legislation_summaries/other/l21069_es.htm [accessed 03.11.14].
39. Marinovich M, Galli CL, Bosetti C, Gallus S, La Vecchia C. Aspartame, low-calorie sweeteners and disease: regulatory safety and epidemiological issues. *Food Chem Toxicol.* 2013;60:109–15.