



SPANISH ASSOCIATION OF PAEDIATRICS

## Criteria for hospital discharge of the healthy term newborn after delivery<sup>☆</sup>



Segundo Rite Gracia<sup>a,\*</sup>, Alejandro Pérez Muñuzuri<sup>b</sup>, Ester Sanz López<sup>c</sup>, José Luis Leante Castellanos<sup>d</sup>, Isabel Benavente Fernández<sup>e</sup>, César W. Ruiz Campillo<sup>f</sup>, M. Dolores Sánchez Redondo<sup>g</sup>, Manuel Sánchez Luna<sup>c</sup>, en representación del Comité de Estándares, de la Sociedad Española de Neonatología

<sup>a</sup> Hospital Universitario Miguel Servet, Zaragoza, Spain

<sup>b</sup> Hospital Clínico Universitario, Santiago de Compostela, A Coruña, Spain

<sup>c</sup> Hospital Universitario Gregorio Marañón, Madrid, Spain

<sup>d</sup> Hospital Universitario Santa Lucía, Cartagena, Murcia, Spain

<sup>e</sup> Hospital Universitario Puerta del Mar, Cádiz, Spain

<sup>f</sup> Hospital Universitario Vall d'Hebron, Barcelona, Spain

<sup>g</sup> Complejo Hospitalario de Toledo, Toledo, Spain

Received 11 August 2016; accepted 22 August 2016

Available online 3 April 2017

### KEYWORDS

Newborn;  
Discharge;  
Early neonatal  
discharge

**Abstract** Criteria for newborn hospital discharge have to include physiological stability and family competence to provide newborn care at home. In this document, the Committee of Standards of the Spanish Society of Neonatology reviews the minimum criteria to be met before hospital discharge of a term newborn infant. We include a review of hospital discharge criteria for the late preterm infants, as these infants are often not hospitalised and remain with their mother after birth. A shortened hospital stay (less than 48 h after delivery) for healthy term newborns can be considered, but it is not appropriate for every mother and newborn. Newborn infants discharged before 48 h of age, should be examined within 3–4 days of life.

© 2016 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. All rights reserved.

### PALABRAS CLAVE

Recién nacido;  
Alta;  
Alta neonatal precoz

### Criterios de alta hospitalaria del recién nacido a término sano tras el parto

**Resumen** Los criterios para el alta de un recién nacido deben incluir la estabilidad fisiológica y la competencia de la familia para proporcionar los cuidados al recién nacido en el domicilio.

<sup>☆</sup> Please cite this article as: Rite Gracia S, Pérez Muñuzuri A, Sanz López E, Leante Castellanos JL, Benavente Fernández I, Ruiz Campillo CW, et al. Criterios de alta hospitalaria del recién nacido a término sano tras el parto. An Pediatr (Barc). 2017;86:289.

\* Corresponding author.

E-mail address: [sriteg@salud.aragon.es](mailto:sriteg@salud.aragon.es) (S. Rite Gracia).

En este documento, el Comité de Estándares de la Sociedad Española de Neonatología revisa los criterios de mínimos que se deben dar antes del alta de un recién nacido a término. Se incluye una revisión de los criterios de alta en el caso de recién nacidos prematuros tardíos, debido a que estos recién nacidos frecuentemente no son hospitalizados y permanecen con sus madres tras el nacimiento. Se puede considerar, en recién nacidos a término sanos, una estancia hospitalaria reducida (menor a 48 h tras el nacimiento), pero esta no es apropiada para todas las madres y todos los recién nacidos. Aquellos recién nacidos dados de alta antes de las 48 h del nacimiento deben ser evaluados entre el tercer y el cuarto día de vida.

© 2016 Asociación Española de Pediatría. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

## Introduction

The hospital stay of the mother and newborn must be long enough to allow the identification of problems and guarantee that the mother is recovered and prepared enough to care for herself and the newborn at home. Cardiopulmonary problems related to the transition from the intrauterine to the extrauterine environment usually become apparent within 12 h from birth. However, other problems such as jaundice, ductal-related cardiac lesions, gastrointestinal obstruction, etc., may require a longer period of observation.<sup>1</sup> The length of stay of the mother–infant dyad after delivery has progressively decreased in the past few decades.<sup>2</sup> At present, the usual length of stay for healthy newborns in Spain is of at least 48 h for vaginal deliveries and 72–96 h for caesarean deliveries. This practice is recommended, as in general, existing diseases may manifest during this period, while it also allows for a more accurate assessment of initial feedings and bowel movements and for the time to provide the care and recommendations needed for transitioning to the care of the newborn at home.<sup>3</sup>

The criteria for NB discharge must include physiologic stability, family preparedness and competence to provide adequate care for the NB at home, and guaranteed access to the health care system and resources. An inadequate assessment in any of these areas prior to discharge can put the NB at increased risk and may result in readmission. Several epidemiological studies have used readmission rates to assess the adequacy of the NB hospital length of stay. However, they have reported readmissions after an early discharge ranging from no increase to a significant increase.<sup>2,4,5</sup> These studies consistently identified jaundice, dehydration, feeding difficulties and infection as the most common reason for readmission.<sup>6,7</sup> Other frequently reported risk factors for readmission are primiparity, maternal comorbidity, shorter gestation, lower birth weight, assisted delivery and small size for gestational age.<sup>1</sup>

**Readiness for discharge.** Perceptions of readiness or unreadiness of the family often differ among paediatricians, midwives, and the mothers themselves. Factors associated with perceptions of unreadiness for newborn discharge include first live birth, maternal history of chronic disease or illness after birth, neonatal illness in the early hours

**Table 1** Grade for recommendations for specific clinical preventive action.

<b>A</b> There is good evidence to recommend the clinical preventive action
<b>B</b> There is fair evidence to recommend the clinical preventive action
<b>C</b> The existing evidence is conflicting and does not allow to make a recommendation for or against use of the clinical preventive action; however, other factors may influence decision-making
<b>D</b> There is fair evidence to recommend against the clinical preventive action
<b>E</b> There is good evidence to recommend against the clinical preventive action
<b>I</b> There is insufficient evidence to make a recommendation; however, other factors may influence decision-making

Source: Canadian Task Force on Preventive Health Care.<sup>9</sup>

post birth, feeding difficulty, history of inadequate prenatal care and poor social support. In light of this, the American Academy of Paediatrics has recommended the use of checklists that can aid clinicians with the preparation of a newborn for discharge.<sup>1,8</sup> The toolkit proposed by the American Academy of Paediatrics mainly focuses on the risk of severe hyperbilirubinaemia, access to breastfeeding support, and coordination of care to improve care for newborns following discharge.

All efforts should be made to keep mothers and newborns together to promote simultaneous discharge.

In this review, we analysed the criteria for the discharge of healthy term newborns, as well as specific criteria for late preterm newborns, who at present often do not require admission to inpatient care and can remain with their mothers. Furthermore, we will analyse the requisites that must be fulfilled for the safe early discharge of newborns.

The grade for recommendations was based on the criteria established by the Canadian Task Force on Preventive Health Care (Table 1).<sup>9</sup>

## Recommendations for the discharge of healthy term newborns

The length of stay of a healthy term newborn must be determined based on the health of the mother, the health and stability of the newborn, the capacity and confidence of the mother in providing care for the newborn, the support available at home and access to appropriate follow-up care. All efforts must be made to keep mothers and newborns together to promote breastfeeding on demand and simultaneous discharge.

Some pathological processes can have an early onset (first 24–48 h post birth), even if the newborn seemed healthy at the time of birth and in the first hours of life. Thus, it is recommended that health providers specially trained in paediatrics conduct an evaluation prior to discharge from the hospital.<sup>3</sup>

The establishment of adequate feeding patterns in the first hours of life is essential to prevent the development of hypernatraemic dehydration problems, which occur frequently when feeding is insufficient.<sup>10,11</sup> The subsets of healthy term newborns at highest risk of dehydration are those delivered by caesarean section and those that are exclusively breastfed.<sup>12</sup> Since the length of stay is short, in some cases there is not enough time to verify that feeding is well established, so the weight of the newborn must be measured to compare it to the weight recorded soon after birth. This allows the rigorous monitoring of newborns with losses of more than 10% of their birth weights.<sup>13</sup>

As we have already mentioned, jaundice is one of the most frequent reasons for readmission.<sup>6,7</sup> The risk of readmission for jaundice depends on several factors, including the length of stay, and lower rates of early discharge have been associated with lower rates of readmission due to jaundice.<sup>14,15</sup> The findings of the medical examination along with the total serum bilirubin measurement can help predict the risk of subsequent severe bilirubinaemia,<sup>16</sup> and this prediction is highly accurate when the pre-discharge total serum bilirubin is combined with gestational age.<sup>17</sup> Furthermore, there is evidence that the implementation of pre-discharge jaundice screening programmes with the use of appropriate nomograms reduces the risk of subsequent admission for significant hyperbilirubinaemia.<sup>18</sup> While exposure to natural light may reduce bilirubin levels, its use in our setting is not justified due to the risks associated with sun exposure of a skin surface area large enough to be effective, and therefore is recommended against. There is no evidence that direct exposure, exposure through a pane of glass or exposure to environmental light have beneficial effects in the prevention or treatment of neonatal jaundice.

It is recommended that the following minimum criteria be met before discharge of a term newborn following an uncomplicated gestation and delivery:

1. Healthy term newborns should not be discharged before 48 h from birth if delivered vaginally or 72–96 h if delivered by caesarean section (B).
2. Absence of evidence of abnormalities in the newborn during the hospital stay and the routine physical examination. This examination must be performed by a physician with a paediatrics specialty at least once, or by a physician with documented experience in neonatal care and under the supervision of a paediatrics specialist (B).
3. It is recommended that the weight, hydration and nutritional status of the newborn be assessed prior to discharge, especially in exclusively-breastfed newborns or those delivered by caesarean section (B).
4. Normal and stable vital signs for at least 12 h prior to discharge; axillary temperature between 36.5 and 37.4 °C, respiratory rate below 60 bpm and no other signs of respiratory distress and an awake heart rate of 100–160 bpm.<sup>19–21</sup> A resting heart rate of up to 70 bpm with no signs of circulatory compromise and adequate responsiveness is also acceptable. A heart rate close to or above the upper bound of the normal range requires further assessment (B).
5. The newborn has urinated regularly and passed at least one stool spontaneously (B).
6. The newborn has completed at least two successful feedings, with assessment to verify that the newborn is able to coordinate sucking, swallowing and breathing while feeding (B).
7. Routine bilirubin measurement in all newborns is not indicated. If the newborn presents with significant jaundice before discharge or is discharged before 48 h (early newborn discharge) the total serum bilirubin levels must be measured and compared with risk charts to determine whether the newborn needs phototherapy. Under these circumstances, an appropriate follow-up plan must be instituted (B). Newborns at risk of significant hyperbilirubinaemia must be identified and assessed<sup>22</sup>:
  - a. Gestational age of less than 38 weeks.
  - b. A previous sibling with neonatal jaundice requiring phototherapy.
  - c. Mother's intention to breastfeed exclusively.
  - d. Visible jaundice in the first 24 h of life.
8. The risk factors for infection have been assessed, and, when present, the newborn has been evaluated appropriately and according to established guidelines for the management of newborns with suspected early-onset bacterial sepsis<sup>23</sup> (A).
9. The results of maternal blood tests, newborn blood type and direct Coombs test have been reviewed<sup>1</sup> (B).
10. Due to the change in the vaccination calendar and delay in the first dose of the Hepatitis B vaccine, a high level of cover must ensure a high coverage for the prenatal screening of pregnant women, with vaccination and immune globulin prophylaxis of newborns of HBsAg+ mothers in the first 24 h post birth<sup>24</sup> (A).
11. Newborn metabolic<sup>25</sup> and hearing<sup>26</sup> screenings have been completed according to the specific protocols established for each (A).
12. The mother has been given information and education on how to provide adequate care for her infant,<sup>1</sup> and the acquisition of this knowledge and competencies has been confirmed (B).
13. Instructions have been given in regard to the subsequent followup of the newborn, emphatically recommending a first visit in primary care within 72 h of discharge, which ideally would be scheduled between the hospital and

primary care prior to discharge.<sup>3</sup> Directions to follow in case of a complication or emergency must also be given (B).

14. It is advisable to ensure that there have been no misinterpretations of the provided information on the part of the family due to language barriers or physical, psychological or sensory impairments<sup>3</sup> (B).
15. Social risk factors have been assessed.<sup>1</sup> When any are identified, discharge should be delayed until they are resolved or a plan has been developed to guarantee the safety of the newborn (B).

## Recommendations for the discharge of late preterm newborns

Late preterm newborns, with a gestational age of 34<sup>+0</sup> to 36<sup>+6</sup> weeks at birth, frequently receive care in level I units and remain with their mothers after delivery. However, they are immature from a physiological and metabolic standpoint. Consequently, they are at higher risk than term newborns of developing complications, which are associated with increased morbidity and mortality, and have higher rates of readmission during the neonatal period, most frequently due to jaundice, feeding difficulties, dehydration and sepsis.<sup>27</sup>

The risk factors for readmission or neonatal morbidity identified in this population include: first born child, exclusive breastfeeding at discharge and history of maternal complications during delivery.<sup>28,29</sup>

The criteria and recommendations for the discharge of late preterm newborns are similar to those for healthy term newborns. However, given the increased risk of neonatal morbidity, the parents of late preterm newborns need to be given specific instructions before discharge and require a more thorough followup thereafter. It is important to educate parents on how to assess for adequate feeding and which signs to watch for to detect dehydration and hyperbilirubinaemia at an early stage. In some instances, this education may require a longer stay after birth. In 2014, the SEN34-36 group of the Spanish Society of Neonatology (Sociedad Española de Neonatología) developed guidelines for the perinatal management and followup of late preterm newborns.<sup>30</sup> We would like to highlight the recommendations pertaining to discharge in this article.

The minimum criteria for discharge are similar to those for healthy term newborns, although the following points must be emphasised:

1. Gestational age has been calculated through appropriate methods.
2. The length of stay after birth must be determined on a case-to-case basis and be based on feeding ability, adequate thermoregulation and the absence of disease and social risk factors. Late preterm newborns may not have developed the abilities required for discharge before 48 h post birth (B).
3. Medical followup in primary care 24–48 h after discharge must be arranged prior to discharge (B). The SEN34-36 group of the Spanish Society of Neonatology

recommends a schedule of weekly checkups until 40 weeks' postmenstrual age.

4. Adequate breastfeeding or bottle feeding for a minimum of 24 h has been verified, with evidence of adequate coordination, suction and breathing during feedings (B). The SEN34-36 of the Spanish Society of Neonatology recommends that weight loss during the hospital stay do not exceed 7% of the birth weight.
5. Feeding technique has been assessed thoroughly during the hospital stay following birth (B).
6. A feeding plan has been established, and the family understands it (B).
7. The risk of developing severe hyperbilirubinaemia has been assessed. As noted above, combining the findings of the medical examination with the total serum bilirubin measurement helps predict the risk of severe hyperbilirubinaemia. In this regard, measurement of total serum bilirubin prior to discharge is recommended in all late preterm newborns, especially those that are exclusively breastfed (B).
8. The physical examination of the newborn was conducted by a physician with a paediatrics specialty and ruled out anomalies that could require a longer stay (B).
9. Potential social risk factors have been assessed. When such factors are identified, discharge should be delayed until they are resolved or social services become involved (B).
10. The mother and other potential caregivers have received sufficient information and education to provide adequate care to the infant after discharge, with particular emphasis on specific issues pertaining to late preterm newborns (B).
11. Information on the prevention of sudden infant death has also been provided (A).

## Early newborn discharge

In recent decades, the length of stay after delivery has been declining steadily from eight to ten days in 1950 to two to four days today in the countries in our region, changing the concept of early newborn discharge over time.<sup>31</sup> Thus, at present an early newborn discharge corresponds to discharge between 24 and 48 h post birth, always in newborns from uncomplicated gestations and deliveries. Very early newborn discharge is defined a discharge before 24 h post birth.

Some of the potential risks associated with early discharge are those derived from the failure to detect certain anomalies, hypernatraemic dehydration secondary to insufficient feeding, and severe jaundice. However, there are no controlled studies demonstrating an increased risk of readmission, and only one retrospective study conducted in Canada.<sup>32</sup> This study compared early discharges in the 1989–1990 and The 1996–1997 periods, and found an inverse correlation between length of stay in days and the risk of readmission. However, the authors did not analyse whether any type of postpartum care was provided at home or in primary care settings in the early days of life.

Some of the advantages of early newborn discharge include increased parental satisfaction and that it

facilitates mother–child bonding and the establishment of breastfeeding.<sup>33</sup> In this sense, the decision to discharge must be determined on a case-to-case basis taking into account the mother’s perception of her ability to take on the care of the newborn and herself. Another advantage would be the optimisation of health care resources resulting from early discharge. Cost analyses performed in Spain have demonstrated cost cuts of 18–20% compared to the usual lengths of stay.<sup>34</sup>

One essential feature of early discharge programmes is the establishment of a postpartum care plan for the mother and infant that ideally would include a home visit service and guaranteeing at minimum one paediatric visit at the hospital or primary care centre at three to four days post birth.<sup>1,15,31,35,36</sup> In this regard, one study found that a group of infants enrolled in an early discharge programme that included home visits made fewer emergency department visits in the first three months of life compared to the group that was discharged following the usual protocol without home visits.<sup>37</sup> On the other hand, another two studies did not find differences between home visits and office visits in the associated rates of readmission, duration of breastfeeding and incidence of postpartum depression.<sup>38,39</sup>

Recommendations for early newborn discharge:

1. Discharge of the newborn before 48 h post birth is possible if the following criteria are met (B):
  - a. Absence of family-related, social or environmental risk (teen mother, mental illness, history of domestic violence during pregnancy, substance use, lack of family support or inability of the mother to care for the NB).
  - b. Records of maternal blood tests (HIV, hepatitis B and syphilis).
  - c. Born to term with a weight appropriate for gestational age.
  - d. Normal medical examination findings and vital signs. Absence of disease requiring admission.
  - e. Absence of risk of infection requiring observation.
  - f. Adequate thermoregulation.
  - g. Ability to feed orally and to appropriately coordinate sucking, swallowing and breathing.
  - h. Urination and meconium defecation (at least one bowel movement before discharge).
  - i. In case the maternal blood group is O or Rh-negative, verification that the direct Coombs test of the newborn is negative.
  - j. Absence of significant hyperbilirubinaemia.
  - k. A sample has been collected for metabolic screening. In addition to the sample before discharge, collection of a second sample is recommended at 5–7 days of life.
    - l. Hearing screening has been performed.
2. A written protocol on early newborn discharge must be in place (B).
3. The capacity of the mother to adequately care for the newborn must be assessed before making the decision to discharge (B).
4. The family will be provided with a report covering not only the care given during the stay, but also the post-discharge care and follow-up plan (B).
5. It must be possible for the newborn to have a postnatal checkup (at a hospital or primary care clinic) at 3 or 4

days of life, and this visit should be scheduled prior to discharge from the maternity ward (B).

Should discharge before 24 h post birth be requested (very early newborn discharge), the parents will be informed of the risks of discharge under these circumstances and asked to sign an informed consent form.

## Conclusions

The timing of newborn discharge must be decided by a paediatric-specialty physician on the basis of strictly clinical criteria and involving the family in the decision-making process. A shortened hospital stay (of less than 48 h following birth) can be considered for healthy term newborns; however, this is not an option for every mother and newborn. In addition to meeting a series of criteria before discharge, early newborn discharge requires a postnatal paediatric visit at day 3 or 4 post birth, which can take place at the hospital or primary care centre. The establishment of standardised processes such as the use of pre-discharge checklists may facilitate a more uniform implementation of the recommendations in the different health facilities that provide neonatal care throughout Spain.

## Conflicts of interest

The authors have no conflicts of interest to declare

## References

1. Benitz WE, Committee on Fetus and Newborn. Hospital stay for healthy term newborns infants. *Pediatrics*. 2015;135:948–53.
2. Datar A, Sood N. Impact of postpartum hospital-stay legislation on newborn length of stay: readmission, and mortality in California. *Pediatrics*. 2006;118:63–72.
3. Sánchez Luna M, Pallás Alonso CR, Botet Mussons F, Echániz Urcelay I, Castro Conde JR, Narbona E. Recomendaciones para el cuidado y atención al recién nacido sano en el parto y en las primeras horas después del nacimiento. *An Pediatr (Barc)*. 2009;71:349–61.
4. Watt S, Sword W, Krueger P. Longer postpartum hospitalization options – who stays, who leaves, what changes? *BMC Pregnancy Childbirth*. 2005;5:13.
5. Paul IM, Lehman EB, Hollenbeak CS, Maisels MJ. Preventable newborn readmissions since passage of the Newborn’s and Mother’s Health Protection Act. *Pediatrics*. 2006;118:2349–58.
6. Escobar GJ, Greene JD, Hulac P, Kincannon E, Bischoff K, Gardner MN, et al. Rehospitalisation after birth hospitalisation: patterns among infants of all gestations. *Arch Dis Child*. 2005;90:125–31.
7. Danielsen B, Castles AG, Damberg CL, Gould JB. Newborn discharge timing and readmissions: California, 1992–1995. *Pediatrics*. 2000;106 1 Pt 1:31–9.
8. Academia Americana de Pediatría. Safe and healthy beginnings: a resource toolkit for hospitals and physician’s offices. 2009 [consulted 3 Jul 2016]. Available in: <https://www.aap.org/en-us/presionarios-resources/quality-improvement/Quality-Improvement-Innovation-Networks/Pages/Safe-and-Healthy-Beginnings-A-Resource-Toolkit-for-Hospitals-and-Physicians-Offices.aspx?nfstatus=401>.

9. Canadian Task Force on Preventive Health Care. New grades for recommendations from the Canadian Task Force on Preventive Health Care. *CMAJ*. 2003;169:207–8.
10. Martínez Escribano MP, Sierra Sirvent J, Mengual Gil J, Membrado Granizo J, Loris Pablo C. Deshidratación hipernatrémica con lactancia materna. *An Esp Pediatr*. 1984;21:774–6.
11. Unal S, Arhan E, Kara N, Uncu N, Aliefendioglu D. Breast-feeding-associated hyponatremia: retrospective analysis of 169 term newborns. *Pediatr Int*. 2008;50:29–34.
12. Konetzny G, Bucher HU, Arlettaz R. Prevention of hypernatraemic dehydration in breastfed newborn infants by daily weighing. *Eur J Pediatr*. 2009;168:815–8.
13. Uras N, Karadag A, Dogan G, Tonbul A, Tatli MM. Moderate hyponatremic dehydration in newborn infants: retrospective evaluation of 64 cases. *J Matern Fetal Neonatal Med*. 2007;20:449–52.
14. Burgos AE, Schmitt SK, Stevenson DK, Phibbs CS. Readmission for neonatal jaundice in California, 1991–2000: trends and implications. *Pediatrics*. 2008;121:e864–9.
15. Meara E, Kotagal UR, Atherton HD, Lieu TA. Impact of early newborn discharge legislation and early follow-up visits on infant outcomes in a state Medicaid population. *Pediatrics*. 2004;113:1619–27.
16. Newman TB, Liljestrand P, Escobar GJ. Combining clinical risk factors with serum bilirubin levels to predict hyperbilirubinemia in newborns. *Arch Pediatr Adolesc Med*. 2005;159:113–9.
17. Keren R, Luan X, Friedman S, Saddlemire S, Cnaan A, Bhutani VK. A comparison of alternative risk-assessment strategies for predicting significant neonatal hyperbilirubinemia in term and near-term infants. *Pediatrics*. 2008;121:e170–9.
18. Eggert LD, Wiedmeier SE, Wilson J, Christensen RD. The effect of instituting a prehospital-discharge newborn bilirubin screening program in an 18-hospital health system. *Pediatrics*. 2006;117:e855–62.
19. Mayfield SR, Bathia J, Nakamura KT, Rios GR, Bell EF. Temperature measurement in term and preterm neonates. *J Pediatr*. 1984;104:271–5.
20. Taylor WC, Watkins GM. Respiratory rate patterns in the newborn infant. *Can Med Assoc J*. 1960;83:1292–5.
21. Semizel E, Oztürk B, Bostan OM, Cil E, Ediz B. The effect of age and gender on the electrocardiogram in children. *Cardiol Young*. 2008;18:26–40.
22. National Institute for Health and Care Excellence. Jaundice in newborn babies under 28 days. NICE guidelines [CG98]. Published date: May 2010. Last updated: May 2016 [consulted 3 Jul 2016]. Available in: <https://www.nice.org.uk/guidance/cg98>.
23. Polin RA, Papile LA, Baley JE, Benitz W, Carlos W, Cummings J, et al., Committee on Fetus and Newborn. Management of neonates with suspected or proven early-onset bacterial sepsis. *Pediatrics*. 2012;129:1006–15.
24. Barker JD Jr, Anuras S, Summers R. Gamma globulin prophylaxis of viral hepatitis. An update. *Postgrad Med*. 1976;60:90–4.
25. American Academy of Pediatrics Newborn Screening Authoring Committee. Newborn screening expands: recommendations for paediatricians and medical homes – implications for the system. *Pediatrics*. 2008;121:192–217.
26. American Academy of Pediatrics, Joint Committee on Infant Hearing. Year 2007 position statement: principles and guidelines for early hearing detection and intervention programs. *Pediatrics*. 2007;120:898–921.
27. Engle WA, Tomashek KM, Wallman C, Committee on fetus and newborn, American academy of pediatrics. “Late-preterm” infants: a population at risk. *Pediatrics*. 2007;120:1390–401.
28. Shapiro-Mendoza CK, Tomashek KM, Kotelchuck M, Barfield W, Weiss J, Evans S. Risk factors for neonatal morbidity and mortality among “healthy”, late preterm newborns. *Semin Perinatol*. 2006;30:54–60.
29. Tomashek KM, Shapiro-Mendoza CK, Weiss J, Kotelchuck M, Barfield W, Evans S, et al. Early discharge among late preterm and term newborns and risk of neonatal mortality. *Semin Perinatol*. 2006;30:61–8.
30. Hurtado-Suazo JA, García-Reymundo M, Calvo-Aguilar MJ, Ginovart-Galiana G, Jiménez-Moya A, Trincado-Aguinagalde MJ, et al. Recomendaciones para el manejo perinatal y seguimiento del recién nacido prematuro tardío. *An Pediatr (Barc)*. 2014;81:327.e1–7.
31. Cuidados desde el nacimiento. Recomendaciones basadas en pruebas y buenas prácticas. Madrid: Ministerio de Sanidad y Política Social; 2010.
32. Liu S, Wen SW, McMillan D, Trouton K, Fowler D, McCourt C. Increased neonatal readmission rate associated with decreased length of hospital stay at birth in Canada. *Can J Public Health*. 2000;91:46–50.
33. Romero J, Romero J, Picazo B, Tapia L, Díaz R, Romero I. Alta hospitalaria temprana en recién nacidos a término sanos. *An Esp Pediatr*. 1999;50:479–84.
34. Sainz J, Ruiz M, Garrido R, Gutiérrez A, Fernández A, Almeida C, et al. Early discharge from obstetrics–pediatrics at the Hospital de Valme, with domiciliary follow-up. *Am J Obstet Gynecol*. 2005;193 3 Pt 1:714–26.
35. Escobar GJ, Braveman PA, Ackerson L, Odouli R, Coleman-Phox K, Capra AM, et al. A randomized comparison of home visits and hospital-based group follow-up visits after early postpartum discharge. *Pediatrics*. 2001;108:719–27.
36. Nelson VR. The effect of newborn early discharge follow-up program on pediatric urgent care utilization. *J Pediatr Health Care*. 1999;13:58–61.
37. Cooper W, Kotagal U, Atherton H, Lippert C, Bragg E, Donovan E. Use of health care services by inner-city infants in an early discharge program. *Pediatrics*. 1996;98:686–91.
38. Lieu TA, Braveman PA, Escobar GJ, Fischer AF, Jensvold NG, Capra AM. A randomized comparison of home and clinic follow-up visits after early postpartum hospital discharge. *Pediatrics*. 2000;105:1058–65.
39. Gagnon A, Dougherty G, Jimenez V, Leduc N. Randomized trial of postpartum care after hospital discharge. *Pediatrics*. 2002;109:1074–80.